

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 773 294 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
17.09.1997 Bulletin 1997/38

(51) Int. Cl.⁶: **C12N 15/16**, C07K 14/71,
C12N 15/10, C12N 15/85

(43) Date of publication A2:
14.05.1997 Bulletin 1997/20

(21) Application number: 96115534.8

(22) Date of filing: 30.07.1990

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI LU NL SE

(30) Priority: 28.07.1989 US 386053

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
90912327.5 / 0 484 440

(71) Applicant: THE UNITED STATES OF AMERICA
as represented by the Secretary
UNITED STATES DEPARTMENT OF COMMERC
E
Washington, DC 20231 (US)

(72) Inventors:


- Miki, Toru
Rockville, MD 20852 (US)
- Aaronson, Stuart A.
Great Falls, VA 22066 (US)
- Fleming, Timothy
Derwood, MD 20855 (US)

(74) Representative: Killin, Stephen James et al
Venner, Shipley & Co.,
20 Little Britain
London EC1A 7DH (GB)

(54) Efficient directional genetic cloning system

(57) A highly efficient genetic cloning system is disclosed which is particularly useful for cloning cDNA copies of eukaryotic mRNAs and can direct the orientation of inserts in plasmid composite vectors with large cloning capacities. Cleavage of such vector DNA, by the restriction enzyme SfiI, for example, creates two different non-symmetrical 3' extensions at the ends of vector DNA. Using a linker-primer and an adaptor, cDNA is prepared to have two different sticky ends which can be ligated to those of the vector. When the cDNA fragments and the vector DNAs are mixed, both the molecules can assemble without self-circularization due to base-pairing specificity. This system provides (1) high cloning efficiency (10^7 - 10^8 clones/g poly (A)⁺ RNA), (2) low background (more than 90% of the clones contain inserts), (3) directional insertion of cDNA fragments into the vectors, (4) presence of a single insert in each clone, (5) accommodation of long inserts (up to 10kb), (6) a mechanism for rescue of the plasmid part from a λ genome, and (7) a straightforward protocol for library preparation.

Figure 15a



EP 0 773 294 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 11 5534

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	SCIENCE, vol. 245, 1989, US, pages 57-60, XP002026815 PAULINE L. LEE ET AL.: "Purification and complementary DNA cloning of a receptor for fibroblast growth factor." * the whole document * -----	1	C12N15/16 C07K14/71 C12N15/10 C12N15/85
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07K
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
THE HAGUE		3 March 1997	DELANGHE, L
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 (12.92) (P04.001)



European Patent Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

See sheet -B-

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☒ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims: 1-10,18,19



European Patent Office

EP 96 11 5534 - B -

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions, or groups of inventions, namely:

1. Claims 1-10,18,19 : DNA encoding a keratinocyte growth factor receptor, vector containing DNA, host, protein and process.
2. Claims 11-17 : Method of identifying the presence in a DNA of phenotypically identifiable trait.

12

EUROPEAN PATENT APPLICATION

21 Application number: **87301031.8**

51 Int. Cl.⁴: **C 12 N 9/22**

22 Date of filing: **05.02.87**

C 12 N 9/96

// C12N15/00

30 Priority: **05.02.86 US 826220**

43 Date of publication of application:
02.09.87 Bulletin 87/36

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **Szybalski, Wacław**
1124 Merrill Springs Road
Madison Wisconsin 53705 (US)

72 Inventor: **Szybalski, Wacław**
1124 Merrill Springs Road
Madison Wisconsin 53705 (US)

74 Representative: **Bass, John Henton et al**
REDDIE & GROSE 16 Theobalds Road
London WC1X 8PL (GB)

54 **Universal restriction endonuclease.**

57 A universal restriction endonuclease for cleaving a target DNA at a predetermined site. The universal restriction endonuclease utilizes a tailored oligodeoxynucleotide adapter in conjunction with a restriction enzyme which cuts DNA at a known distance from its recognition site. The oligodeoxynucleotide adapter mimics the relationship which exists in such enzymes between its recognition site and cut site.

Descripti n

UNIVERSAL RESTRICTION ENDONUCLEASEBackground of the Invention

5 The present invention relates to novel universal restriction endonucleases and to the methods of producing such endonucleases. More specifically, the present invention relates to restriction endonucleases which allow one to cleave practically any predetermined DNA sequence by combining an oligodeoxynucleotide adapter with a known restriction endonuclease.

10 Restriction endonucleases are a class of enzymes that occur naturally in bacteria. When they are purified away from other interfering bacterial components, restriction endonucleases can be used in the laboratory to cut DNA molecules into precise fragments. This property enables DNA molecules to be uniquely identified and to be precisely fractionated into their constituent subunits. Restriction endonucleases have proved to be indispensable tools in modern genetic research. They are the biochemical 'scissors' by means of which genetic engineering and analysis is performed.

15 Restriction endonucleases are classified into three groups. Class-I and Class-III enzymes carry modification (methylation) and an ATP-requiring restriction (cleavage) activity in the same protein moiety. Both types of enzymes recognize unmethylated recognition sequences in substrate DNA, but Class-I enzymes cleave largely randomly, whereas Class-III enzymes cut DNA at specific sites, generally at a constant distance from their recognition sites.

20 Class-II restriction/modification systems consist of a separate restriction endonuclease and modification methylase. These enzymes cut DNA within or near their particular recognition sequences, which typically are four to six nucleotides in length, frequently with a twofold axis of symmetry. There exist more than 500 Class II restriction endonucleases that cleave DNA at specific sites. The number of different recognition sites is greater than 100 (not counting various modifications, including methylation) (Roberts, Nucl. Acids Res., 13:165, 1985; Kessler et al. Gene, 33:1-102, 1985). Although the existing repertoire of cleavage sites is of great value, there is frequently a need for additional cut sites to permit precise engineering of genes and regulatory sequences. At present, to access a sequence where a favorable restriction site is not available, one usually engineers the sequence by a variety of methods, including insertion of appropriate linkers, mutagenesis, or exonucleolytic digestion. However, such methods for creating a new site modify the nucleotide sequence under study and may often affect its function.

SUMMARY OF THE INVENTION

In accordance with the present invention, there is provided a novel approach to the engineering of restriction enzymes which can cleave DNA at a specific predetermined site. In one embodiment Class IIS or "shifter" enzymes are utilized. Class IIS enzymes are defined as those Class II endonucleases which cleave double-stranded (ds) DNA at precise distances from recognition sequences. Certain Class III enzymes may also be used in practicing the present invention, particularly those which cut at a reasonably constant distance (at about 20-25 nucleotides) from their recognition site. Class II enzymes with cut sites between two recognition sites (e.g., BstXI enzyme; see Gene 40:173 (1985)) may also be used in accordance with the teachings of the present invention.

40 The present invention utilizes the separation between the recognition site and the cut site to allow Class II or Class III enzymes to cleave practically any predetermined sequence by combining the enzyme with a properly designed oligodeoxynucleotide adapter. The adapter is constructed from the constant recognition site domain (a hairpin-containing ds sequence) and a variable, single-stranded (ss) domain complementary to the target ss sequence to be cleaved. The ss sequence to be cleaved may be provided by ss phage DNA, gapped ds plasmids, supercoiled ds plasmids that are alkali denatured and rapidly neutralized, or the like. Combination of these three components, i.e. the Class IIS enzyme, the ss DNA target sequence, and the complementing adapter, result in a novel restriction-endonuclease complex which cleaves the target DNA at a specific predetermined site. A further advantage of the present invention is that the target ss DNA may be converted to the precisely cleaved ds DNA by an appropriate DNA polymerase, utilizing the adapter oligodeoxynucleotide as a primer.

50 Thus, in accordance with the present invention, there is provided a method of changing the enzyme specificity of restriction endonucleases by synthetic design leading to a practically unlimited assortment of new endonucleases having predetermined restriction specificities.

BRIEF DESCRIPTION OF THE DRAWINGS

55 Figure 1 illustrates the design scheme of an oligodeoxynucleotide adapter for instructing the FokI Class IIS enzyme to cleave M13 ss DNA at a predetermined point.

Figure 2 illustrates two oligodeoxynucleotide adapters for use with the FokI restriction endonuclease.

60 Figure 3 shows a schematic representation of the M13mp7 ss DNA.

Figure 4 shows reproduction of photographs of gel digests of ds and ss M13mp7 DNA by FokI, HphI, HpaI, MboI, MnI, SfaNI and BbyI.

Figure 5 illustrates conformation of an oligodeoxynucleotide adapter and its insusceptibility to FokI

cleavage.

Figure 6 illustrates photographs of gels of FokI digestion of M13mp7 in the presence of the 32-mer and 34-mer adapters illustrated in Fig. 2.

Figure 7 illustrates photographs of gels of a preformed FokI enzyme-adapter complex used to digest M13mp7. The effects of salt concentrations and annealing conditions between the adapter and target DNA are also shown.

Figure 8 shows a schematic representation of M13mp7 ds DNA.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides purposefully designed oligodeoxynucleotide adapters which can be used to modify the specificity of restriction endonucleases. The adapters instruct these enzymes to cleave DNA at a predesignated target site. In other words, the oligodeoxynucleotide adapters, which may be likened to novel synthetic "coenzymes", confer a new substrate specificity upon a restriction endonuclease.

The oligodeoxynucleotide adapters of the present invention are used to mimic the relationship which exists in certain restriction endonucleases between their recognition sites and cut sites so as to provide the modified endonuclease with the ability to cut DNA at practically any predetermined site on a target DNA. Preferred restriction enzymes, modified in accordance with the present invention, are class IIS enzymes for which the cleavage site is shifted a considerable (up to 13 nucleotides), but precise, distance from the recognition site. The recognition sites and displacements of cleavage sites of 11 Class IIS restriction enzymes are listed in Table I. In accordance with the present invention, such Class IIS enzymes may be engineered by using synthetic oligodeoxynucleotide adapter molecules to instruct the enzyme to recognize and cut any predetermined DNA sequence. As noted above, Class III enzymes, preferably those which cut at about 20-25 nucleotides from their recognition site, and other Class II enzymes, e.g., BstXI, may also be modified to function as a universal restriction endonuclease by the use of analogous adapters in accordance with the teachings of the present invention. Such enzymes include EcoPI, EcoP15, HinI and HinIII; see Kessler et al., supra, pp.67-68.

TABLE I
Properties of class IIS restriction endonucleases ("shifters")

Enzyme	Recognition sequences	Protruding end	Reference	
			Page	Entry
BbvI ^{d,e}	5'-GCAGCN ⁸ CGTCGN ¹²	5'-N ⁴	57	513
BbvII	5'-GAAGACN ² CTTCTGN ⁶	5'-N ⁴	102	-
BlnI	5'-GGATCN ⁴ CCTAGN ⁵	5'-N ¹	57	516
FokI ^{d,e}	5'-GGATGN ⁹ CCTACN ¹³	5'-N ⁴	57	519
HgaI ^{d,f}	5'-GACGCN ⁵ CTGCGN ¹⁰	5'-N ⁵	58	526
HphI ^{d,e}	5'-GGTGAN ⁸ CCACTN ⁷	3'-N ¹	58	527
MboI ^{d,e}	5'-GAAGAN ⁸ CTTCN ⁷	3'-N ¹	58	528

TABLE I (continued)
Properties of class IIS restriction endonucleases ("shifters")

Enzyme	Recognition Sequences	Protruding end	Reference Page	Entry
<u>Nnl</u> Id,f	5'-CCTCN7 GGAGN7	5'-N7GAGG N7CTCC	blunt	59 531
<u>Sfa</u> Id,f	5'-GCATCN5 CTGAGN9	5'-N9GATGC N5CTACG	5'N4	59 532
<u>Taq</u> II	G G 5'-CACCCAN11 CTGGCTN9 G G	G G 5'-N9 TCGGTC N11ACCCAC G G	3'N2	59 533
<u>Tth</u> IIIIII9	5'CAAPuCAN11 GTPPyGTN9	5'N9 TGPpyTTG N11ACPuAAC	3'N2	59 535

- a. Since the recognition sites are not palindromic, both orientations are listed. Arrows specify the cleavage sites shown only for the first enzyme.
- b. The 5' or 3' protruding ss ends of the restriction sites are indicated and their lengths given.
- c. Refers to the review of Kessler et al., (supra), with the page and entry number specified.
- d. Commercially available (e.g., New England Biolabs; see Kessler et al., supra).
- e. No endonucleolytic activity toward M13 ss DNA.
- f. The tested batch of the enzyme displayed an endonucleolytic activity toward M13 ss DNA. However, the enzymes may be purified to remove the endonucleolytic activity.
- g. Py= T or C; Pu= A or G. For cut points see Roberts, supra.

Referring now to Fig. 1, there is shown the outline of the principle of the present invention as applied to the FokI endonuclease. The relationship between the FokI recognition site, GGATC, CCTAC, and the staggered

sites on the normal ds DNA molecule are shown in Fig. 1a. The adapter oligodeoxynucleotide (Fig. 1b) that mimics the above arrangement consists of the hairpin ds domain, which carries the FokI recognition sequence (underlined), and an ss domain where the cleavage(s) would occur after pairing with the complementary target DNA strand. Thus, the ds domain of the adapter is a constant part that provides the specificity of the Class II enzyme, whereas the ss domain is a variable and can be designed in accordance with the present invention to provide cleavage at any chosen site in the complementary target DNA.

DESIGN OF THE ADAPTER

The oligodeoxynucleotide depicted in Fig. 1b will serve as an example of adapter design for the Class II FokI enzyme and ss target DNA from the M13 phage. The hairpin ds domain of the 34-mer contains the 5'-GGATC recognition site for FokI; its ss domain 3'-CCTAC was designed to direct the cleavage between T and C in the target M13 DNA. The complex between the adapter and M13 DNA (Fig. 1c) mimics a ds module (Fig. 1a) which contains both the FokI recognition site and the ds cleavage site (see Table I and Fig. 1c). Depending on the sequence of the ss domain of the adapter, the enzyme would cut at any chosen site on the target DNA. At the same time, the 3' end of the adapter (see Fig. 1b,c) could serve as primer for synthesis of the strand complementary to the target ss DNA, either in one stage using all four dNTPs, or in two stages: a limited synthesis with three dNTP's (to anchor the 3' end of the adapter before FokI digestion, but not so extensive as to provide ds recognition sites for FokI), followed by synthesis of the total complementary strand after the FokI cleavage.

Fig. 1 does not indicate optimal adapter design. Optimization includes: (i) determination of the most efficient hairpin lengths to the left and to the right of the GGATC recognition site; (ii) alteration of the length of the ss domain complementary to the cleaved ss DNA and/or to provide the primer function; (iii) modification of the position of the ss M13 DNA branch (Fig. 1b) with respect to the GGATC and cleavage sites; (iv) comparison of adapters with the 5' and 3' ss domains (see Fig. 2); (v) possible use of ambiguous nucleotides (inosinic acid) or ribonucleotides in the ss domain of the adapter; and (vi) inclusion of reactive deoxynucleotides (e.g., 5-iododeoxyuridine nucleotides) in the tip of the hairpin in order to form covalent (but active) adapter-enzyme complexes upon light UV irradiation. Other methods of adapter-enzyme complexing could also be used. Optimization of adapter design also facilitates a better understanding of the mechanism of the endonucleolytic activity of restriction endonucleases.

Although it is necessary to design a specific adapter for each chosen cleavage site, only the variable ss domain (5-15 nucleotides) has to be custom-made. It could then be either (i) ligated to the stem of the constant ds hairpin domain, using as a splicing support a set of oligodeoxynucleotides designed to be complementary to the adjoining sequences of the hairpin and ss domains or, (ii) prepared by first synthesizing a supply of the constant hairpin domain with protective groups retained, and in the second synthetic stage adding the variable ss domain. Such two-step organic synthesis is possible for adapters containing the 5' ss domain (see Fig. 2a), because of the 3' to 5' direction of the nucleotide coupling.

Although adapter synthesis must be customized for many sequences, in accordance with the present invention, an inventory of adapters for large numbers of commonly encountered sequences for which no enzymes are presently available may be prepared.

TARGET DNA AND VECTORS

(1) Single-stranded DNA

One of the main advantages of using ss target-vector DNA, as exemplified by M13 ss DNA shown in Fig. 1, is that it does not provide recognition sites for most of the Class IIS enzymes, which are rather frequent cutters of ds vectors. Several commercially available Class IIS enzymes (FokI, HphI, MboI, BbvI; New England Biolabs) do not have interfering endonucleolytic activity toward M13 DNA. The ss target DNA could be obtained from a variety of sources, including the ss DNA phage vectors (M13, fd, fl) or any ds plasmid carrying the replication origin (ori) of those phages that convert ds plasmids to packaged ss DNA upon phage superinfection.

(2) Double-stranded DNA with ss gaps

Another substrate could be ds DNA with properly designed ss gaps, but in this case the ds cleavage sites should preferably be either absent or rendered inactive by (i) methylation or other modification of the ds DNA, or (ii) saturation of enzyme with the adapter and possible covalent binding. For the specific methylation of FokI sites, a FokI methylase is available (New England Biolabs). For class IIS enzymes which create 4-nucleotide cohesive ends, the religation of the cleaved plasmid would restore its original arrangement, since each cleavage sequence is unique. However, a cut in the ss gap would remain open, if the uncut adapter were removed. If the purpose of the adapter-mediated FokI cleavage is to produce a small fragment with a specific terminus, the additional FokI cuts in the ds DNA might be of no importance or even beneficial.


(3) Double-stranded DNA

In addition to the problems specified above, use of a ds DNA substrate requires a special method for threading the ss domain of the adapter into the ds target sequence. This should be possible when using reversibly denatured target DNA obtained by alkali denaturation of the supercoiled plasmid DNA followed by rapid neutralization. Such a procedure results in largely denatured molecules shown to anneal with primer

oligonucleotides (Chen and Seeburg, DNA 4:165-170, 1985). Use of a single adapter would result in an ss nick, whereas two adapters with barely overlapping or neighboring ss domains complementary to opposing strands would provide staggered cuts. Alternatively, this might be accomplished by use of catalysts, e.g., RecA-like protein, mild denaturation conditions, or of ribonucleotides in the ss domain of the adapter. Also, advantageously, with the adapter shown in Figs. 1 or 2b, the ds DNA may be obtained by using the 3'-OH end of the adapter as a primer to synthesize the complementary strand, as discussed above.

The present invention permits construction of novel cleavage specificities by combining carefully contrived synthetic adapter molecules with Class IIS or Class III restriction endonucleases. Devising unlimited numbers of new enzymatic specificities ("new restriction enzymes") in accordance with the present invention is limited mainly by oligodeoxynucleotide synthesizer capacity.

Although the present invention is primarily directed to Class IIS restriction endonucleases, it is also possible, in accordance with the teachings herein, to design adapters for enzymes that recognize an inverted repeat separated by a run of nonspecific base pairs. For instance, the adapter



should convert the BstXI enzyme (see Kessler et al., *supra*) to an endonuclease seeking an ss sequence complementary to the central NNNNNN ss domain of the adapter. Many other kinds of adapters could also be designed.


The following examples are given to additionally illustrate embodiments of the present invention as it is presently preferred to practice. It should be understood that these examples are illustrative, and that the invention is not to be considered as restricted thereto except as indicated in the appended claims.

EXAMPLE 1

Example I describes methods designed to select target DNA, target sequence, restriction enzyme, as well as methods to detect cleavage processes.

(a) Selection of the target DNA: For the purposes of this example, a phage M13 derivative was selected as the source of ss DNA because (i) its entire nucleotide sequence is known (Van Wezenbeek et al., Gene 11:129-148 1980; Yanish-Perron et al, Gene 33: 103-119 1985), (ii) M13mp phages are convenient cloning vectors, and (iii) it is easy to propagate both the ds replicative form (RF) and ss DNA. Among M13 derivatives, the M13mp7 vector phage was selected because its ss DNA contains a small hairpin structure which was designed to be cleaved by enzymes EcoRI, BamHI, HindIII, AccI and SalI (Messing, Methods in Enzymol. 101:20-78, 1983). These cuts provided reference points for confirmatory mapping of the ss cuts by the FokI-adapter complexes, as depicted in Fig. 3.

The heavy lines of Fig. 3 represent the ss DNA. The sequence of the hairpin "a" is as follows:



Two cuts at position I (Fig.3) are as described in Fig. 2. The location of cut(s) at position II was found to be located near nucleotide 6938. The length of the entire M13mp7 is 7238 nucleotides, and the approximate lengths of the fragments are: a, 32; b, 670; c, 1640; d, 4890; b + c, 2300; a + b + d, 5600 nucleotides. The drawing is not to scale.

(b) Selection of the target sequences: Using the computer program of the University of Wisconsin Genetics Computer Group on a DESCS VAX 7800 (Devereux et al., Nucleic Acids Res. 12:387-375 1984), a 14-nucleotide sequence in M13mp7 ss DNA was found which satisfied the following criteria: (i) it is unique, (ii) no similar sequences with one, two or three mismatches are present in the phage DNA, (iii) no complementary sequences with 0-2 mismatches are present in the phage, and (iv) the cuts in the selected sequence and at the reference EcoRI site (see cuts I and EcoRI in Fig. 3) would produce two fragments easy to distinguish and measure on gels (ignoring the 32-nucleotide hairpin fragment cut-off by EcoRI). A 14-nucleotide unique sequence 5'-TGCTACCCTCGTTC-3' was selected (nucleotides 1339-1352), with no 0-3 nucleotide mismatches found in M13mp7 ss DNA, with only three 4-nucleotide scattered mismatches (starting at nucleotide 1749, 2908 and 5599), and no 0-2 nucleotide-mismatch-complementary sequences, which might have interfered with the adapter pairing. The sole 3-nucleotide mismatch complementary sequence starts at nucleotide 5443.

(c) Selection of the restriction enzyme: Testing of all the commercially available Class IIS enzymes revealed that four of them (FokI, HphI, MboI and BbvI) do not show endonucleolytic activity toward ss M13mp7 (Fig. 4).

Fig. 4 illustrates cleavage of the replicative form RF ds (lanes 1,3,5,7,9,11,13) and ss (lanes 2,4,6,8,10,12,14) M13mp7 DNA by *FokI* (8 u; lane 1,2), *HphI* (8 u; lanes 3,4), *HpaI* (2 u; lanes 5,6) *Mb* *II* (20 u; lanes 7,8), *MnII* (2 u; lanes 9,10), *SfaNI* (2 u; lanes 11,12) and *BbvI* (2 u; lanes 13,14). All enzymes were a gift from New England Biolabs and digestions were carried out for 3 hours at 37°C under conditions suggested by the manufacturer for the ds DNA, using the indicated number of enzyme units (u; see above in parentheses) per 20 µl of digestion mix per 2 µg DNA. Conditions of sample preparation and electrophoresis of ss M13mp7 DNA were the same as described below in connection with Fig. 6. Conditions for sample preparation with digestion of ds DNAs were as in Maniatis et al. (1982). DNA was stained with ethidium bromide and photographed using UV light trans-illumination. For the first phase *FokI* was selected because of previous experience with this enzyme and because a *FokI* methylase has become commercially available, permitting protection of ds DNA from *FokI* digestion at preexisting restriction sites.

As seen in Fig. 4, Class IIS enzymes *HgaI* and *MnII* show multiple cuts on M13mp7 ss DNA, whereas *SfaNI* seems to contain some nonspecific endonucleolytic activity toward ss DNA. All seven enzymes show the predicted ds cuts, with the exception of *MbolI*, where Dam methylation interferes with a few cuts (see also Kessler et al., *supra*).

(d) Design of the adapters: Since the *FokI* recognition sequence

GGATG
CCTAC

is asymmetric, two different oligodeoxynucleotide adapters, "left" (32 nucleotides) and "right" (34 nucleotides), were constructed, as shown and explained in Fig. 2. They were expected to direct two different cuts, 8 nucleotides apart, on the M13mp7 target ss DNA (Figs. 2 and 3). The adapter is composed of two domains (see Fig. 1): (i) the constant ds (hairpin) domain which contains the *FokI*-recognition ds site, and (ii) the variable ss domain complementary to the target sequence described in section b above. The left adapter, with the hairpin ds domain at its 3' end, permits its two-stage synthesis: (i) first, the 3'- to 5'-organic synthesis of a large supply of the constant ds domain in the protected form, and (ii) subsequent addition of several predesigned ss domains complementary to the selected target sequence. The right adapter, with the hairpin ds domain at its 5' end, permits the use of its 3' end as the primer for synthesis of the entire complementary strand of M13mp7 DNA.

(e) Conformation of the adapters: As seen in Fig. 5, the adapter could exist as a hairpin monomer (see Fig. 2) or form a dimer, as e.g. that corresponding to the right adapter:

5'-CACATCCGTGCACGGATGTGGAACGAGGGTAGCA-3'
3'-ACGATGGGAGCAAGGTGTAGGCACGTGCCTACAC-5'

with the central 20-base pair palindrome susceptible to several restriction enzymes, including *HgiAI* and *Bsp1286*. When electrophoresis was carried out under non-denaturing conditions (Fig. 5, A-lane 1 and B-lane 1) the dimers were observed. When the alkaline sample mix or 70°C treatment in the loading buffer were employed, the dimer bands disappeared (Fig. 5B, lanes 3 and 2, respectively). In the *FokI* buffer the 34-mer denaturation at 70°C was partially reversed (Fig. 5A, lane 3). Both monomer and dimer should be active as adapters.

(f) Sensitivity of the adapters to the *FokI* enzyme: Since it was desired that the *FokI* enzyme cut only the duplex formed by annealing between the M13 target sequence and the adapter, it was necessary to demonstrate that the enzyme would not cleave the ss domain of the adapter in the absence of complementary DNA strand. Fig. 5 illustrates conformation of the adapter and its insusceptibility to *FokI* cleavage.

In Fig. 5a, all lanes contain 1 µg of 34-nucleotide adapter suspended in *FokI* buffer (see Fig. 6) and heated for 5 min at 70°C (lanes 3-4) or thermally untreated (lanes 1-2). Adapter in lanes 2 and 4 was treated with *FokI*, as described in connection with Fig. 6. Electrophoresis was performed on 20% PA gel in presence of Peacock buffer (Maniatis et al., 1982) at 200 volts, for 5 h. Each well was filled in with 20 µl of adapter solution and 5 µl of loading buffer containing bromophenol blue (Maniatis et al., 1982).

In Fig. 5b, all lanes contain 2 µg of 32-nucleotide adapter suspended either in 20 µl of loading buffer (10% glycerol, 7% sucrose, 2.4% Ficoll; free of bromophenol blue) - lanes 1 and 2; or in 20 µl of alkaline loading buffer - lane 3. Samples in lane 2 were heated 5 min at 70°C. Samples (20 µl) were electrophoresed as in panel A. Bands were stained with ethidium bromide and photographed as in Fig. 4 (a) dimer, (b) monomer, (c) bromophenol blue indicator. This proved to be the case as shown in Fig. 5A (lanes 2 and 4), since adapter alone is not cleaved by *FokI*. As mentioned in section (d) above, the dimerization of the adapter molecules was observed.

(g) Detection of cleavages in ss M13mp7 DNA: Agarose gel (1.4%) electrophoresis was used to detect cleavage of the ss circular DNA, as described by Been and Champoux, *Methods in Enzymol.* 101:90-98, 1983. Single cuts convert the circles to linear DNA, which migrates more rapidly than the closed circles (Fig. 6; conversion from CF to LF). Digestion with *EcoRI* served as a control, since there is an *EcoRI* site in the hairpin formed by the invert repeat in M13mp7 DNA (Fig. 3).

EXAMPLE II

Example II describes the use of the enzyme-adapter complex to cleave ss DNA at a predetermined site.

(a) Digestion of M13mp7 ss circular DNA with the *FokI* restriction enzyme in presence of the left (32-mer) and right (34-mer) adapters: Figure 6 illustrates electrophoretic separation of linear and circular forms of M13mp7 ss DNA carried out in horizontal 1.4% agarose gels in Tris-borate (Peacock) buffer for 3 h at 120 V,

and generally following the procedure of Been and Champoux, *supra*. M13mp7 ss circular DNA was prepared as described by Maniatis et al. (1982). Each lane contains 2 μ g of this DNA (i) suspended in total volume of 20 μ l of FokI buffer (20 mM KCl, 10 mM Tris. HCl, pH 7.5, 10 mM MgCl₂, 0.5 mM DTT), (ii) digested or otherwise treated, (iii) precipitated with 4 vols. of 95% ethanol in presence of 0.3 M Na acetate, for 10 min. at -70°C, and centrifuged in microfuge for 10 min, (iv) washed with 70% ethanol and dried under vacuum, and resuspended in 20 μ l of alkaline sample mix (30 mM NaOH, 2 mM EDTA, 7% Ficoll, 0.1% SDS, and 0.01% bromphenol blue). DNA bands were stained with ethidium bromide (see Fig. 4).

Lane 1 (Fig.6) contains untreated M13mp7 DNA [over 90% of circular form (CF), upper band, and less than 10% of linear form (LF), lower band]. Lane 2 contains the same DNA treated with 12 units FokI/20 μ l for 2 h at 37°C in FokI buffer (see above). Lanes 3 and 4 contain the same DNA hybridized with 15-fold molar excess of 32-mer and 34-mer adapter, respectively. Hybridization procedure consisted of 5 min. heating at 70°C followed by 1 h at 37°C in FokI buffer. There is no DNA cleavage in lanes 1-4, which represent negative controls. Lanes 5 and 6 represent the same treatment as in lanes 3 and 4 followed by FokI digestions as in lane 2. Here one sees specific cleavage of M13 ss DNA. Lane 7 represents, as a positive control, i.e., digestion with EcoRI (16 u/20 μ l, 1 h, 27°C) in FokI buffer plus Tris•HCl to 100 mM (pH 7.5). Lanes 8 and 9 represent the same treatment as in lanes 5 and 6 followed by 1 h EcoRI digestion as in lane 10 which allows location of FokI adapter cut sites. Designations a+b+d and c refer to bands in lanes 5 and 6 of Fig. 6 and to the corresponding fragments in Fig. 3. For size markers see Fig. 7.

(a) As shown in Fig. 6 (lanes 1-4), neither FokI alone nor the adapters alone cleave circular M13mp7 DNA. However, when either the left or right adapter was first annealed to the ss DNA followed by addition of FokI, the circles were linearized with over 90% efficiency. Both adapters participate in the endonucleolytic reaction, but the right (34-nucleotide) adapter seems to be more efficient.

(b) Secondary cuts by the FokI-adapter combination: Careful examination of lanes 5 and 6 in Fig. 6 and lanes 2-6 in Fig. 7 shows two more bands in addition to the 7238-nucleotide band representing the entire linear ss DNA of M13mp7. The appearance of the two secondary bands seems to depend on the conditions of the reaction.

(c) Mapping of the cuts in M13mp7 DNA: To determine the positions of the primary and secondary cuts, M13mp7 ss DNA was cleaved with both the FokI-adapter complex and the reference EcoRI enzyme, which cuts within the M13mp7 hairpin as shown in Fig. 3. The results shown in lanes 8 and 9 (Fig. 6) appear to be consistent with the positions of the FokI cuts as shown in Fig. 3. Cleavages at site I are located within the 14-nucleotide target sequences (nucleotides 6938-6951; Fig. 2), and cleavage(s) at site II seems to occur at a perfect 9-nucleotide match (nucleotides 1340-1348) between the adapter and the M13 ss DNA as shown by nucleotide sequencing and indicated by the dots:

```

6938          6951
5'-AGCTACCCTCTCCG-3'
      .....
      TGCTACCCTCGTTC
1339          1352

```

The upward arrow specifies the cut in the ss DNA directed by the 34-nucleotide adapter. Cleavage directed by the left 32-nucleotide adapter should be on the right edge of the 9-nucleotide homology (Fig. 2), consistent with the lower yield of secondary cuts (compare lanes 5 and 6, Fig. 6). (The last digits of the numbers match the corresponding nucleotides).

EXAMPLE III

Example III illustrates various conditions of the cleavage reaction.

(a) Cleavage with the preformed enzyme-adapter complex: In the procedure described in Example II, the adapter was annealed to M13mp7 DNA before addition of FokI. Here the enzyme was first saturated with the adapter and only then allowed to react with the target M13mp7 ss DNA.

Thus, when adapter is in excess, the free adapter molecules should bind to the M13 ss target DNA and interfere with cutting by the enzyme-adapter complexes. On the other hand, when enzyme is in excess, such competition should not occur since free enzyme does not recognize the ss target DNA.

Therefore, increasing concentrations of the FokI and a constant amount of adapter to preform the enzyme-adapter complexes were used as follows:

Lane 1 in Fig. 7A corresponds to the untreated M13 circular ss DNA control (as in lane 1, Fig. 6). For lanes 2-4, various amounts of FokI (12, 24, and 36 units, respectively) were combined with 34-mer (amount identical to that in Fig. 5; denatured for 5 min at 70°C) and incubated for 10 min. at 37°C in 20 μ l of FokI buffer. To each enzyme adapter preparation, 2 μ g of M13mp7 circular ss DNA was added and the digestion carried for 1 h at 37°C in the total volume of 30 μ l. Other procedures were as in Fig. 6.

(b) Effect of buffer, KCl and Mg++ concentrations: Lanes 1 and 2 of Fig. 7B correspond to lanes 1 and 6 in Fig. 6. Lanes 3-6 correspond to the following variants of the experiment in lane 2: in lanes 3 and 4, the FokI buffer was diluted twofold and fourfold, respectively, both for hybridization and digestion. In lanes 5 and 6, all

KCl was removed from the *FokI* buffer. Moreover, in lane 6 the Mg^{++} concentration was reduced fourfold (from 10 to 2.5 mM $MgCl_2$).

(c) Effect of the time period of the DNA-adapter hybridization: Lanes 1 and 2 of Fig. 7C correspond to lanes 1 and 2 in panel B. Lanes 3-5 correspond to lane 2, but the time of M13 ss DNA-adapter (34-mer) hybridization was reduced to 30, 15 and 5 min., respectively. Lane 6 (ss size standards) represents ss DNA fragments of the plasmid pRHP233 denatured and electrophoresed in the alkaline sample mix. The size of the fragments (from the top) are: 3636, 3097, 2459, 2443 nucleotides, followed by two doublets (complementary strands) of 1700 and 1159 nucleotides, respectively. Panels (A), (B), and (C) represent separate experiments run on different days.

As shown in Fig. 7A, such complexes do cleave M13 ss DNA, and as predicted above, the cleavage is more efficient at increasing *FokI*/adapter ratios (lanes 2-4). It also appears that use of the preformed enzyme-adapter complex has little effect on the proportion of the secondary cuts (cut II in Fig. 3).

All the reactions discussed in Examples II and III were carried out in buffer consisting of 20 mM KCl, 10 mM Tris-HCl, 10 mM $MgCl_2$ and 0.5 mM DTT. More stringent annealing conditions were explored, i.e., at decreasing salt concentrations, to determine whether the proportion of secondary cuts could be lowered, while cleavage efficiency remains substantially unaffected. As shown in Fig. 7B (lanes 3 and 4), twofold or fourfold dilution of the buffer still permits the reaction to proceed, while the ratio of the primary to secondary cuts seems to be slightly increased. Even total elimination of the KCl component results in the selective and efficient cleavage (lane 5). The most selective cleavage was obtained in the absence of KCl and 2.5 mM $MgCl_2$ (lane 6).

A standard time of 1 h annealing at 37°C between adapter and target M13mp7 ss DNA was utilized before adding the *FokI* enzyme. As shown in Fig. 7C, the annealing time could be reduced to 5 min. with hardly any change of the digestion pattern.

EXAMPLE IV

Example IV describes preparation of double-stranded DNA fragments, using the enzyme-adapter complex. The adapter DNA comprising the 5' double-stranded hairpin domain and 3' ss domain (Fig. 1 and Fig. 2b) may be used as a primer for the synthesis of the complementary DNA strand. This adapter was annealed with the ss M13mp7 template, with adapter attaching at sites I and II (Fig. 3). Upon addition of Polk (Klenow fragment of *E. coli* DNA polymerase I) and all four dNTP's, the DNA synthesis was carried out for 0.5-2 hrs at 20°C in the *FokI* buffer. *FokI* enzyme was added either 5 min after initiation of complementary DNA synthesis or at the end of reactions. In both cases two new DNA fragments were produced, and are initiated and cut at site I (1341) and the other at site II (6938) (see Figs. 3 and 8). The sizes of novel fragments were 1116 bp and 603 bp, because they were also cut at the regular *FokI* sites at nucleotide 225 and 6335 (Fig. 8). Digestion of double-stranded RF form of M13mp7 DNA produced only four fragments, 3335, 2775, 910 and 218 bp long (Fig. 8). Thus, the 1116-bp and 603-bp fragments were novel and produced by the novel enzymatic activity of the adapter-*FokI* complex. The 1116-bp fragment was produced with high yield because there is a perfect match between the adapter and M13mp7 as DNA at site I (nucleotide 1341). The yield of the 603-bp fragment was lower because only 9 out of 14 nucleotides of adapter did pair at site II (Figs. 3 and 8); however, Polk was able to use this primer with only 9 bp match, even though the 3'-terminal nucleotide of the adapter was not paired. The resulting novel DNA fragments (with blunt ends due to the Polk reaction) were blunt-end cloned into M13mp7 vector and their ends were sequenced. In general, the sequences matched the predicted sequences of the novel ends. This method of preparing double-stranded DNA fragments with one predetermined terminus appears to be both simple and reliable. The second terminus of the fragment could be either at *FokI* site (Fig. 8), or any other end generated by another restriction enzyme of choice.

Claims

1. A universal restriction endonuclease which cleaves a target DNA at a predetermined site comprising:
 - (a) an oligodeoxynucleotide adapter; and
 - (b) an appropriate restriction enzyme.
2. The universal restriction endonuclease of claim 1, wherein the oligodeoxynucleotide adapter and restriction enzyme are covalently bound.
3. The universal restriction endonuclease of claim 1, wherein the oligodeoxynucleotide adapter mimics the relationship which exists in the restriction enzyme between its recognition site and its cleavage site.
4. The universal restriction endonuclease of claim 1, wherein the restriction enzyme is a Class II enzyme selected from the group of *BbvI*, *BbvII*, *BlnI*, *FokI*, *HgaI*, *HphI*, *MboI*, *MnII*, *SfaNI*, *TagII* or *TthIII*.
5. The universal restriction endonuclease of claim 4, wherein the restriction enzyme is *FokI*.
6. The universal restriction enzyme of claim 4, wherein the oligodeoxynucleotide adapter comprises a double stranded nucleotide sequence corresponding to the recognition site of the restriction enzyme and a single stranded nucleotide sequence complementary to the predetermined site of the target DNA.
7. The universal restriction endonuclease of claim 1, wherein the restriction enzyme is a Class III enzyme selected from the group of *EcoPI*, *EcoP15*, *HinI*, or *HinfIII*.
8. The universal restriction enzyme of claim 7, wherein the oligodeoxynucleotide adapter comprises a

double-stranded nucleotide sequence corresponding to the recognition site of the restriction enzyme and a single-stranded nucleotide sequence complementary to the predetermined site of the target DNA.

9. A method for cleaving a target DNA at a predetermined site comprising contacting the target DNA with a modified restriction endonuclease.

10. The method of claim 9, wherein the modified restriction endonuclease comprises:

- (a) an oligodeoxynucleotide adapter; and
- (b) an appropriate restriction enzyme.

11. The method of claim 10, wherein the oligonucleotide adapter and restriction enzyme are covalently bound.

12. The method of claim 10, wherein the oligonucleotide adapter comprises a double-stranded nucleotide sequence corresponding to the recognition site of the restriction enzyme and a single-stranded nucleotide sequence complementary to the predetermined site of the target DNA.

13. The method of claim 10, wherein the contacting of the modified restriction endonuclease with the target DNA comprises

- (a) complexing the oligodeoxynucleotide adapter to the predetermined site of the target DNA; and
- (b) adding the appropriate restriction enzyme to the DNA-adapter complex.

14. The method of claim 13, wherein the target DNA is single stranded.

15. The method of claim 14, comprising the further step of extending the 3'-OH terminus of the oligodeoxynucleotide adapter along the single-stranded target DNA to form a double-stranded DNA containing the predetermined cut site prior to addition of the appropriate restriction enzyme.

16. The method of claim 10, wherein the oligodeoxynucleotide adapter is complexed with the restriction enzyme and the contacting comprises complexing the enzyme-adapter complex with the target DNA

17. The method of claim 16, wherein the target DNA is single-stranded.

18. The method of claim 17, comprising the further step of extending the 3'-OH terminus of the oligodeoxynucleotide adapter along the single-stranded DNA to form a double-stranded DNA, containing the predetermined cut site which double-stranded DNA is then cut by the appropriate restriction enzyme of the enzyme adapter complex.

0235781




FIG. I

0235781




FIG.2

0235781




FIG. 3




FIG. 8

200587

0235781

FokI		HphI		HgaI		MboI		MnII		SfaNI		BbvI	
1	2	3	4	5	6	7	8	9	10	11	12	13	14




FIG. 4

BEST AVAILABLE COPY

0235781




FIG. 5A

FIG. 5B

BEST AVAILABLE COPY

0235781

0235781




FIG.6

BEST AVAILABLE COPY

AD-000000

0235781

1 2 3 4 1 2 3 4 5 6 1 2 3 4 5 6




FIG. 7C

FIG. 7B

FIG. 7A

BEST AVAILABLE COPY